




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/70, C07H 19/10		A1	(11) International Publication Number: WO 99/09998
			(43) International Publication Date: 4 March 1999 (04.03.99)
(21) International Application Number: PCT/US98/17894		Dallas [US/US]; 109 Haley House Lane, Apex, NC 27502 (US).	
(22) International Filing Date: 28 August 1998 (28.08.98)		(74) Agents: BISWAS, Sorojini, J. et al.; Myers, Bigel, Sibley & Sajovec, P.A., P.O. Box 37428, Raleigh, NC 27627 (US).	
(30) Priority Data: 60/057,064 29 August 1997 (29.08.97) US		(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71) Applicants (for all designated States except US): THE UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL [US/US]; 308 Bynum Hall, Campus Box #4105, Chapel Hill, NC 27599-4105 (US). INSPIRE PHARMACEUTICALS [US/US]; Suite 470, 4222 Emperor Boulevard, Durham, NC 27703 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): BOUCHER, Richard, C., Jr. [US/US]; 735 Gimghoul Road, Chapel Hill, NC 27514 (US). SHAVER, Sammy, Ray [US/US]; 614 Hollow Court, Chapel Hill, NC 27516-9459 (US). PENDERGAST, William [US/US]; 5816 Williamsburg Way, Durham, NC 27713-2636 (US). YERXA, Benjamin [US/US]; 1300 Hathaway Road, Raleigh, NC 27608-1936 (US). RIDEOUT, Janet, L. [US/US]; 3101 Morningside Drive, Raleigh, NC 27607-3018 (US). DOUGHERTY, Robert [US/US]; 105 Hunter Hill Road, Chapel Hill, NC 27516 (US). CROOM,		Published With international search report.	

(54) Title: USE OF URIDINE 5'-DIPHOSPHATE AND ANALOGS THEREOF FOR THE TREATMENT OF LUNG DISEASES



(57) Abstract

Compounds of Formula (I) wherein X_1 and X_2 are each independently either O^- or S^- ; X_3 and X_4 are each independently either $-H$ or $-OH$, with the proviso that X_3 and X_4 are not simultaneously $-H$; R_1 is selected from the group consisting of O , imido, methylene and dihalomethylene; R_2 is selected from the group consisting of H , halo, alkyl, substituted alkyl, alkoxy, nitro and azido; R_3 is selected from the group consisting of H , alkyl, acyl, aryl, and arylalkyl; and R_4 is selected from the group consisting of $-OR'$, $-SR'$, $-NR'$, and $-NR'R''$, wherein R' and R'' are independently selected from the group consisting of H , alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxy, and aryloxy, with the proviso that R' is absent when R_4 is double bonded from an oxygen or sulfur atom to the carbon at the 4-position of the pyrimidine ring, are used in methods of hydrating lung mucus secretions and treating lung disorders such as cystic fibrosis, ventilator-associated pneumonia, chronic bronchitis, chronic obstructive pulmonary disorder and primary ciliary dyskinesia. Pharmaceutical compositions containing compounds of Formula (I), and novel compounds of Formula (I) are also described.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

USE OF URIDINE 5'-DIPHOSPHATE AND ANALOGS THEREOF FOR THE TREATMENT OF LUNG DISEASES

10

Related Applications

This application claims the benefit of filing of U.S. Provisional Application No. 60/057,064, filed August 29, 1997, which application is incorporated herein in its entirety.

15

Government Support

This invention was made with United States government support under grant number # 2PO1 HL32322-11A1 from the National Institutes of Health. The United States government has certain rights in this invention

20

Field of the Invention

This invention relates to methods of treating lung disease, and novel compounds and pharmaceutical compositions useful therefor.

25

Background of the Invention

One therapeutic goal in cystic fibrosis and other pulmonary diseases in which the water content of lung mucus is altered is to hydrate the lung mucus secretions, so that the secretions may thereafter be more easily removed from the lungs by mucociliary action or simple coughing. For example, the use of aerosolized amiloride to hydrate mucus secretions is described in U.S. Patent No. 4,501,729 to Boucher et al. Amiloride appears to block Na^+ reabsorption by airway epithelial cells, and therefore inhibits water absorption from the mucus.

30

While an important breakthrough in providing treatments for cystic fibrosis, a potential problem with amiloride as a therapeutic is its relatively short duration of action.

In certain lung diseases (e.g., cystic fibrosis), several functions of airway epithelia are abnormal, and deficiencies in both Cl^- transport and Na^+ absorption are well documented. See, e.g. Knowles et al., *Science* **221**, 1067 (1983); Knowles et al., *J. Clin. Invest.* **71**, 1410 (1983). Regulation of ion transport is thus thought to have potential therapeutic benefit in lung diseases characterized by abnormalities in epithelial ion transport. Confirmation of the presence of P2Y_2 ($\text{P}_{2\text{U}}$ -purinergic) receptors on the apical surface of human airway epithelial cells raised the possibility that aerosolized nucleotides might be used therapeutically to induce Cl^- secretion in individuals with cystic fibrosis or other airway diseases. Accordingly, a different therapeutic approach for hydrating lung mucus secretions is exemplified by techniques that involve the administration of ATP or UTP, which appear to stimulate chloride secretion from respiratory epithelial cells. See, e.g., U.S. Patent No. 5,292,498 to Boucher.

Existence of a G-protein-coupled receptor that selectively recognizes uridine 5'-diphosphate (UDP) was originally established in studies of a receptor natively expressed by C6-2B rat glioma cells. E. R. Lazarowski et al. *J. Biol. Chem.* **269**, 11830-11836 (1994). The P2Y_6 receptor was recently cloned by K. Chang et al., *J. Biol. Chem.* **270**, 26152-26158 (1995). This receptor was subsequently shown to be selectively activated by UDP, and to be the UDP receptor natively expressed in C6-2B cells. R. A. Nicholas et al., *Mol. Pharmacol.* **50**, 224-229 (1996). The failure to identify this receptor in previous studies of mammalian tissues likely has been a consequence of the lack of availability of potent selective agonists for uridine nucleotide receptors, and the low chemical and metabolic stability of the available nucleotides. It was originally reported that UDP stimulated inositol phosphate accumulation in human airway epithelial cells by low potency activation of the P2Y_2 receptor. E. R. Lazarowski et al. *Br. J. Pharmacol.* **116**, 1619-1627 (1995); H. A. Brown et al., *Mol. Pharmacol.* **40**, 648-655 (1991). However, it has been recently demonstrated that UDP is in fact not an


agonist at the P2Y₂ receptor (Nicholas et al., *supra*) and that the previously observed effect of UDP at P2Y₂ receptors can be explained by the presence of small amounts of contaminating UTP in UDP solutions and/or by conversion of UDP to UTP by cell surface nucleoside diphosphokinase.

- 5 Despite the evidence related to the P2Y₆ receptor and its relationship to UDP, it has heretofore not been recognized that this relationship may be useful in the treatment of airway disease.

Summary of the Invention

- 10 The present inventors have discovered that the P2Y₆ receptor, which selectively recognizes UDP as a potent agonist, also exists in airway tissue. The association of the P2Y₆ receptor with increases in Cl⁻ secretion indicates that UDP and other receptor-selective drugs that derive from this molecule are of therapeutic value in the treatment of a variety of airway diseases. Accordingly, a first aspect
- 15 of the present invention relates to a method of hydrating mucus secretions in the lungs of a subject in need of such treatment. The method comprises administering to the lungs of the subject a compound of Formula I below, or a pharmaceutically acceptable salt thereof (hereinafter referred to as the "active compound"), in an amount effective to hydrate lung mucus secretions:

20



wherein:

- 30 X₁, and X₂ are each independently either O⁻ or S⁻ ;
 X₃ and X₄ are each independently either -H or -OH, with the proviso that X₃ and X₄ are not simultaneously -H;

R₁ is selected from the group consisting of O, imido, methylene, and dihalomethylene (*e.g.*, dichloromethylene, difluoromethylene);

R₂ is selected from the group consisting of H, halo, alkyl, substituted alkyl, alkoxyl, nitro and azido;

5 R₃ is selected from the group consisting of H, alkyl, acyl (including arylacyl), and arylalkyl; and

R₄ is selected from the group consisting of -OR', -SR', NR', and NR'R'', wherein R' and R'' are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxyl, and aryloxy, and with the
10 proviso that R' is absent when R₄ is double bonded from an oxygen or sulfur atom to the carbon at the 4-position of the pyrimidine ring.

The method of the present invention may further comprise the step of concurrently administering amiloride, benzamil, or phenamil to the subject in an amount effective to inhibit the reabsorption of water from lung mucus secretions.

15 The method of the present invention is useful in treating several disorders of the lung, including but not limited to, cystic fibrosis, chronic bronchitis, chronic obstructive pulmonary disorder (COPD), primary ciliary dyskinesia, and ventilator-associated pneumonia (VAP).

A second aspect of the present invention is a pharmaceutical composition
20 containing the active compounds disclosed herein, in an amount effective to hydrate lung mucus secretions, in a pharmaceutically acceptable carrier.

Novel compounds useful in the treatment of lung disorders are a third aspect of the present invention. These compounds have the structure of Formula I as set forth above, with the proviso that such novel compounds do not include the
25 known compounds uridine 5'-diphosphate (or UDP), 2-deoxyuridine 5'-diphosphate (or dUDP), uridine 5'-O-(2-thiodiphosphate) (or UDP-β-S), and 4-mercaptouridine 5'-diphosphate (or 4-mercaptoUDP). Novel compounds of the present invention include, but are not limited to, 3'-deoxyuridine 5'-diphosphate; 5-bromouridine 5'-diphosphate; 5-(1-phenylethynyl)-uridine 5'-diphosphate; 5-methyluridine 5'-
30 diphosphate; 4-hexylthiouridine 5'-diphosphate; 4-methoxyuridine 5'-diphosphate; 4-(N-morpholino)uridine 5'-diphosphate; 4-hexyloxyuridine 5'-diphosphate; N,N-

dimethylcytidine 5'-diphosphate; N-hexylcytidine 5'-diphosphate; and N-cyclopentylcytidine 5'-diphosphate.

A fourth aspect of the present invention is the use of the active compounds described herein for the manufacture of a medicament for the therapeutic hydration of mucus secretions in the lungs of a subject in need of such treatment.

Brief Description of the Drawings

FIG. 1 is a graph showing the time course for the conversion of [^3H]UTP to [^3H]UDP in the presence of 10 units/mL hexokinase (HK). The data is expressed as the percentage of [^3H]uridine triphosphate converted to [^3H]uridine diphosphate with open circles (○) indicating the relative amount of [^3H]uridine diphosphate, while filled-in circles (●) indicate the related amount of [^3H]uridine triphosphate. The data indicate the mean value of one experiment representative of two independent experiments performed with duplicate samples that differed by less than 20%.

FIG. 2 illustrates, by means of three separate HPLC traces, the metabolism of [^3H]UTP and [^3H]UDP by human nasal epithelial cells. Each trace represents the results of experiments in which confluent, polarized human airway epithelial cells were incubated for 20 minutes at 37°C in the presence of 1 μM (0.2 μCi) of [^3H]UTP (**FIG. 2A**), [^3H]UDP (**FIG. 2B**), and [^3H]UDP combined with 100 μM ATP (**FIG. 2C**). The traces shown in **FIGS. 2A, 2B and 2C** are representative of at least three independent experiments performed with duplicates. In each trace, the X-axis of the trace indicates elution time in minutes, while the Y-axis indicates the concentration of [^3H] radioactivity in units of $\text{cpm} \times 10^{-3}$.

FIG. 3 is a schematic representation of the effects of UTP and UDP on [^3H]inositol phosphate formation in relation to either the mucosal or serosal cell surface. **FIG. 3** also illustrates the effects of UTP and UDP on intracellular calcium mobilization in polarized human nasal epithelial cells. Confluent cells were loaded with [^3H]myo-inositol and preincubated with LiCl (**FIG. 3A**), or with Fura-2 (**FIG. 3B**), as described below in Examples 2 and 3. The cells were challenged with 100 μM of either UTP (left-hand pair of data bars) or UDP (right-hand set of data bars), added to either the serosal (open bars) or the mucosal

(filled-in bars) medium. The data in FIG. 3A are shown as the concentration of [^3H]inositol phosphate in units of $\text{cpm} \times 10^{-3}$ and represent the medium (\pm S.E.M.) from three experiments performed with triplicates. The data in FIG. 3B are shown as ΔCa^{2+}_i in units of μM , and represent the medium (\pm S.E.M.) from fourteen

5 individual experiments.

FIG. 4 consists of representative tracings of UDP- and UTP-promoted changes in intracellular Ca^{2+} concentration and Cl^- diffusion potentials in human nasal epithelial cells. The upper tracings show changes in intracellular Ca^{2+} concentration on the serosal (left-hand tracing) and mucosal (right-hand tracing) after the addition of UDP and UTP to the cell surface medium, as indicated. The lower tracings illustrate changes of transepithelial potential difference (ΔTEP) on the serosal (left-hand tracing) and mucosal (right-hand tracing) after the addition of UDP and UTP to the cell surface medium, as indicated. The data are representative of at least eight independent experiments.

15 FIG. 5 is a graph illustrating the concentration-response relationship for mucosal UDP- and UTP-stimulated [^3H]inositol phosphate formation in human nasal epithelial cells. The log-concentration of either UTP (filled-in circles, ●) or UDP (filled-in squares, ■) is indicated on the x-axis of the graph. The concentration of [^3H]inositol phosphates in units of $\text{cpm} \times 10^{-3}\text{M}$ is indicated on the y-axis of the graph. The data represent the mean value (\pm S.E.M.) from three independent experiments performed with triplicate samples.

FIG. 6 is a graph illustrating the effects on mucociliary clearance in sheep after the administration of either a saline control (circles, ●) or UDP (squares, ■). Time in minutes after administration of the compound is indicated on the x-axis of the graph, while percentage retention of mucus is indicated on the y-axis.

25 FIG. 7 is a graph illustrating the effects on tracheal mucus velocity (TMV) in sheep after the administration of either a saline control (open diamond ◇) or UDP (closed circle, ●). Time in minutes after administration of the compound is indicated on the x-axis of the graph, while TMV measured as a percentage of baseline is indicated on the y-axis.


30

Detailed Description of the Invention

The methods and pharmaceutical formulations of the present invention can be used to facilitate (i.e., enhance, speed, assist) the clearance of mucus secretions from the lungs of a subject in need of such treatment for any reason, including (but not limited to) retained secretions arising from airway diseases such as cystic fibrosis, chronic bronchitis, chronic obstructive pulmonary disorder (COPD), ventilator-associated pneumonia (VAP), primary ciliary dyskinesia, asthma, bronchiectasis, post-operative atelectasis (plugging of airways with retained secretions after surgery), and Kartagener's syndrome.

The present invention is concerned primarily with the treatment of human subjects, but may also be employed for the treatment of other mammalian subjects, such as dogs and cats, for veterinary purposes.

The methods of the present invention include the administration of compounds of Formula I, while pharmaceutical compositions of the present invention comprise compounds of Formula I. As used herein, a compound of Formula I is as follows:



wherein:

X_1 , and X_2 are each independently either O^- or S^- ;

X_3 and X_4 are each independently either $-H$ or $-OH$, with the proviso that X_3 and X_4 are not simultaneously $-H$;

R_1 is selected from the group consisting of O , imido, methylene, and dihalomethylene (e.g., dichloromethylene, difluoromethylene);

R_2 is selected from the group consisting of H , halo, alkyl, substituted alkyl, alkoxy, nitro and azido;

R₃ is selected from the group consisting of H, alkyl, acyl (including arylacyl), and arylalkyl; and

R₄ is selected from the group consisting of -OR', -SR', NR', and NR'R'', wherein R' and R'' are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxy, and aryloxy, and with the
5 proviso that R' is absent when R₄ is double bonded from an oxygen or sulfur atom to the carbon at the 4-position of the pyrimidine ring.

As used herein the term "alkyl" refers to C₁₋₁₀ inclusive, linear, branched, or cyclic, saturated or unsaturated (i.e., alkenyl and alkynyl) hydrocarbon chains, including for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl,
10 pentyl, hexyl, octyl, ethenyl, propenyl, butenyl, pentenyl, hexenyl, octenyl, butadienyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, and allenyl groups. As used herein, the term "acyl" refers to an organic acid group wherein the -OH of the carboxyl group has been replaced with another substituent (i.e., as represented by
15 RCO—, wherein R is an alkyl or an aryl group). As such, the term "acyl" specifically includes arylacyl groups. Specific examples of acyl groups include acetyl and benzoyl. As used herein, the term "aryl" refers to 5 and 6-membered hydrocarbon and heterocyclic aromatic rings. Specific examples of aryl groups include but are not limited to cyclopentadienyl, phenyl, furan, thiophene, pyrrole,
20 pyran, pyridine, imidazole, isothiazole, isoxazole, pyrazole, pyrazine, pyrimidine, and the like. The term "alkoxy" as used herein refers to C₁₋₁₀ inclusive, linear, branched, or cyclic, saturated or unsaturated oxo-hydrocarbon chains, including for example methoxy, ethoxy, propoxy, isopropoxy, butoxy, t-butoxy, and pentoxy. The term "aryloxy" as used herein refers to phenyloxy or hexyloxy, and alkyl,
25 halo, or alkoxy substituted phenyloxy or hexyloxy. As used herein, the terms "substituted alkyl" and "substituted aryl" include alkyl and aryl groups, as defined herein, in which one or more atoms or functional groups of the aryl or alkyl group are replaced with another atom or functional group, including for example, halogen, aryl, alkyl, alkoxy, hydroxy, nitro, amino, alkylamino, dialkylamino,
30 sulfate, and mercapto. The terms "halo," "halide," or "halogen" as used herein refer to fluoro, chloro, bromo, and iodo groups.

Compounds illustrative of the compounds of Formula (I) above include:
uridine 5'-diphosphate (also referred to herein as UDP); uridine 5'-O-(2-thiodiphosphate) (also referred to herein as UDP- β -S); 2-deoxyuridine 5'-diphosphate (also referred to herein as dUDP); 3'-deoxyuridine 5'-diphosphate (also referred to
5 herein as 3'-deoxyUDP); 5-bromouridine 5'-diphosphate (also referred to herein as 5-BrUDP); 5-(1-phenylethynyl)-uridine 5'-diphosphate (also referred to herein as 5-(1-phenylethynyl)UDP); 5-methyluridine 5'-diphosphate (also referred to herein as 5-methylUDP); 4-hexylthiouridine 5'-diphosphate (also referred to herein as 4-hexylthioUDP); 4-mercaptopuridine 5'-diphosphate (also referred to herein as 4-
10 mercaptoUDP); 4-methoxyuridine 5'-diphosphate (also referred to herein as 4-methoxyUDP); 4-(N-morpholino)uridine 5'-diphosphate (also referred to herein as 4-(N-morpholino)UDP); 4-hexyloxyuridine 5'-diphosphate (also referred to herein as 4-hexyloxyUDP); N,N-dimethylcytidine 5'-diphosphate (also referred to herein as N,N-dimethylCDP); N-hexylcytidine 5'-diphosphate (also referred to herein as N-
15 hexylCDP); and N-cyclopentylcytidine 5'-diphosphate (also referred to herein as N-cyclopentylCDP). Certain compounds of Formula I (e.g., UDP, dUDP, UDP- β -S, and 4-mercaptopuridine) are known and may be made in accordance with known procedures or variations thereof, which will be apparent to those skilled in the art. For example, the identification and preparation of certain thiophosphate analogues of
20 nucleoside diphosphates (such as UTP- β -S) are set forth in U.S. Patent No. 3,846,402 to Eckstein et al., and in R.S. Goody and F. Eckstein, *J. Am. Chem. Soc.* 93, 6252-6257 (1971). Alternatively, UDP, dUDP, and other analogs thereof are also commercially available from vendors such as Sigma (St. Louis, MO) and Pharmacia (Uppsala, Sweden).

25 Other compounds of Formula I useful in the present invention (e.g., 5-(1-phenylethynyl)UDP; 3'-deoxyUDP, 5-methylUDP; 4-hexylthioUDP; 4-methoxyUDP; 4-hexyloxyUDP; 4-(N-morpholino)UDP; N,N-dimethylCDP; N-hexylCDP; and N-cyclopentylCDP) are novel compounds disclosed for the first time herein, and as such are claimed accordingly in the appended claims.

30 For the sake of simplicity, Formula I herein illustrates uridine diphosphate active compounds in the naturally occurring D configuration, but the present invention also encompasses compounds in the L configuration, and mixtures of

compounds in the D and L configurations, unless otherwise specified. The naturally occurring D configuration is preferred.

The active compounds of Formula (I) may be administered by themselves or in the form of their pharmaceutically acceptable salts, e.g., an alkali metal salt such as sodium or potassium, an alkaline earth metal salt, or an ammonium and tetraalkyl ammonium salt, NX_4^+ (wherein X is a C_{1-4} alkyl group). Pharmaceutically acceptable salts are salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects.

Active compounds of the present invention may optionally be administered in conjunction with other compounds useful in the hydration of lung mucus secretions or useful in the facilitation of the removal of lung mucus secretions. Such compounds (herein referred to as "supplemental compounds") include, but are not limited to, benzamil, phenamil, and amiloride. Amiloride (also known as 3,5-diamino-6-chloro-N-(diaminomethylene)pyrazinecarboxamide), benzamil (also known as 3,5-diamino-6-chloro-N-(benzylaminoaminomethylene)pyrazinecarboxamide) and phenamil (also known as 3,5-diamino-6-chloro-N-(phenylaminoaminomethylene)pyrazinecarboxamide) are known compounds and are disclosed in U.S. Patent No. 3,313,813 to E. Cragoe. The terms "amiloride," "benzamil," and "phenamil," as used herein include the pharmaceutically acceptable salts thereof, such as (but not limited to) amiloride hydrochloride, benzamil hydrochloride or phenamil hydrochloride. Amiloride, benzamil or phenamil used to prepare compositions for the present invention may alternatively be in the form of a pharmaceutically acceptable free base of amiloride, benzamil or phenamil. In one embodiment of the invention, the supplemental compound is concurrently administered with the active compound or compounds of the present invention. As used herein, the word "concurrently" means sufficiently close in time to produce a combined (e.g., additive or synergistic) effect. In other words, concurrently may be defined as simultaneously, or it may be defined as two or more events occurring within a short time period before or after each other.

The active and supplemental compounds described herein may be administered to the lungs of a patient by any suitable means, but are preferably administered by administering an aerosol suspension of respirable particles comprised

of the active compound, which the subject inhales. The active compound can be aerosolized in a variety of forms, such as, but not limited to, dry powder inhalants, metered dose inhalants, or liquid/liquid suspensions. The respirable particles may be liquid or solid. The quantity of active compound included may be an amount
5 sufficient to achieve dissolved concentrations of active compound on the airway surfaces of the subject of from about 10^{-9} to about 10^{-1} Moles/liter, and more preferably from about 10^{-6} to about 10^{-4} Moles/liter.

The particulate pharmaceutical composition may optionally be combined with a carrier to aid in dispersion or transport. A suitable carrier such as a sugar (i.e.,
10 lactose, sucrose, trehalose, mannitol) may be blended with the active compound or compounds in any suitable ratio (e.g., a 1 to 1 ratio by weight).

Solid or liquid particulate forms of the active compound prepared for practicing the present invention should include particles of respirable size: that is, particles of a size sufficiently small to pass through the mouth and larynx upon
15 inhalation and into the bronchi and alveoli of the lungs. In general, particles ranging from about 1 to 10 microns in size are within the respirable range. Particles of non-respirable size which are included in the aerosol tend to be deposited in the throat and swallowed, and the quantity of non-respirable particles in the aerosol is preferably minimized.

20 The dosage of active compound will vary depending on the condition being treated and the state of the subject, but generally may be an amount sufficient to achieve dissolved concentrations of active compound on the airway surfaces of the subject of from about 10^{-9} to about 10^{-1} Moles/liter, and more preferably from about 10^{-6} to about 10^{-4} Moles/liter. Depending upon the solubility of the particular
25 formulation of active compound administered, the daily dose may be divided among one or several unit dose administrations. The daily dose by weight will depend upon the age and condition of the subject. Such a daily dose may be as low as 1 mg per day, under certain circumstances may be as low as 0.5 mg per day, and may even be as low as 0.1 mg/day. The daily dose of the active compounds may also be as high as
30 200 mg/day, under certain conditions may be as high as 500 mg/day, and may even be as high as 1000 mg/day. The doses of the active compounds may be provided as one or several prepackaged units.

In the manufacture of a formulation according to the invention, active compounds of the present invention or the pharmaceutically acceptable salts or free bases thereof are typically admixed with, inter alia, an acceptable carrier. The carrier must, of course, be acceptable in the sense of being compatible with any other
5 ingredients in the formulation and must not be deleterious to the patient. The carrier may be a solid or a liquid, or both, and is preferably formulated with the compound as a unit-dose formulation which may contain from 0.5% to 99% by weight of the active compound. One or more active compounds may be incorporated in the formulations of the invention, which formulations may be prepared by any of the well-known
10 techniques of pharmacy consisting essentially of admixing the components.

Aerosols of liquid particles comprising the active compound may be produced by any suitable means, such as with a pressure-driven aerosol nebulizer or an ultrasonic nebulizer. *See, e.g.*, U.S. Patent No. 4,501,729. Nebulizers are commercially available devices which transform solutions or suspensions of the
15 active ingredient into a therapeutic aerosol mist either by means of acceleration of compressed gas, typically air or oxygen, through a narrow venturi orifice or by means of ultrasonic agitation. Suitable formulations for use in nebulizers consist of the active ingredient in a liquid carrier, the active ingredient comprising up to 40% w/w of the formulation, but preferably less than 20% w/w. The carrier is typically water
20 (and most preferably sterile, pyrogen-free water) or a dilute aqueous alcoholic solution, preferably made isotonic but may be hypertonic with body fluids by the addition of, for example, sodium chloride. Optional additives include preservatives if the formulation is not made sterile, for example, methyl hydroxybenzoate, antioxidants, flavoring agents, volatile oils, buffering agents and surfactants.

25 Aerosols of solid particles comprising the active compound may likewise be produced with any solid particulate medicament aerosol generator. Aerosol generators for administering solid particulate medicaments to a subject produce particles which are respirable, as explained above, and generate a volume of aerosol containing a predetermined metered dose of a medicament at a rate suitable for
30 human administration. One illustrative type of solid particulate aerosol generator is an insufflator. Suitable formulations for administration by insufflation include finely comminuted powders which may be delivered by means of an insufflator or taken

into the nasal cavity in the manner of a snuff. In the insufflator, the powder (e.g., a metered dose thereof effective to carry out the treatments described herein) is contained in capsules or cartridges, typically made of gelatin or plastic, which are either pierced or opened *in situ* and the powder delivered by air drawn through the device upon inhalation or by means of a manually-operated pump. The powder employed in the insufflator consists either solely of the active ingredient or of a powder blend comprising the active ingredient, a suitable powder diluent, such as lactose, and an optional surfactant. The active ingredient typically comprises from 0.1 to 100 w/w of the formulation. A second type of illustrative aerosol generator comprises a metered dose inhaler. Metered dose inhalers are pressurized aerosol dispensers, typically containing a suspension or solution formulation of the active ingredient in a liquified propellant. During use these devices discharge the formulation through a valve adapted to deliver a metered volume, typically from 10 to 200 μ l, to produce a fine particle spray containing the active ingredient. Suitable propellants include certain chlorofluorocarbon compounds, for example, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane and mixtures thereof. The formulation may additionally contain one or more co-solvents, for example, ethanol, surfactants, such as oleic acid or sorbitan trioleate, antioxidants and suitable flavoring agents.

Any propellant may be used in carrying out the present invention, including both chlorofluorocarbon-containing propellants and non-chlorofluorocarbon-containing propellants. Thus, fluorocarbon aerosol propellants that may be employed in carrying out the present invention including fluorocarbon propellants in which all hydrogens are replaced with fluorine, chlorofluorocarbon propellants in which all hydrogens are replaced with chlorine and at least one fluorine, hydrogen-containing fluorocarbon propellants, and hydrogen-containing chlorofluorocarbon propellants. Examples of such propellants include, but are not limited to: $\text{CF}_3\text{-CHF-CF}_2\text{H}$; $\text{CF}_3\text{-CH}_2\text{-CF}_2\text{H}$; $\text{CF}_3\text{-CHF-CF}_3$; $\text{CF}_3\text{-CH}_2\text{-CF}_3$; $\text{CF}_3\text{-CHCl-CF}_2\text{Cl}$; $\text{CF}_3\text{-CHCl-CF}_3$; $\text{cy-C(CF}_2)_3\text{-CHCl}$; $\text{CF}_3\text{-CHCl-CH}_2\text{Cl}$; $\text{CF}_3\text{-CHF-CF}_2\text{Cl}$; $\text{CF}_3\text{-CHCl-CFHCl}$; $\text{CF}_3\text{-CFCl-CFHCl}$; $\text{CF}_3\text{-CF}_2\text{-CF}_2\text{H}$; $\text{CF}_3\text{-CF}_2\text{-CH}_3$; $\text{CF}_2\text{H-CF}_2\text{-CFH}_2$; $\text{CF}_3\text{-CF}_2\text{-CFH}_2$; $\text{CF}_3\text{-CF}_2\text{-CH}_2\text{Cl}$; $\text{CF}_2\text{H-CF}_2\text{-CH}_3$; $\text{CF}_2\text{H-CF}_2\text{-CH}_2\text{Cl}$; $\text{CF}_3\text{-CF}_2\text{-CF}_2\text{-CH}_3$; $\text{CF}_3\text{-CF}_2\text{-CF}_2\text{-CF}_2\text{H}$; $\text{CF}_3\text{-CHF-CHF-CF}_3$; $\text{CF}_3\text{-O-CF}_3$; $\text{CF}_3\text{-O-CF}_2\text{H}$; $\text{CF}_2\text{H-H-O-CF}_2\text{H}$; $\text{CF}_2\text{H-O-CFH}_2$;

CF₃-O-CH₃; CF₃-O-CF₂-CF₂H; CF₃-O-CF₂-O-CF₃; cy-CF₂-CF₂-O-CF₂-; cy-CHF-CF₂-O-CF₂-; cy-CH₂-CF₂-O-CF₂-; cy-CF₂-O-CF₂-O-CF₂-; CF₃-O-CF₂-Br; CF₂H-O-CF₂-Br; and mixtures thereof, where "cy" denotes a cyclic compound in which the end terminal covalent bonds of the structures shown are the same so that the end
5 terminal groups are covalently bonded together. Particularly preferred are hydrofluoroalkanes such as 1,1,1,2-tetrafluoroethane (propellant 134a) and heptafluoropropane (propellant 227). A stabilizer such as a fluoropolymer may optionally be included in formulations of fluorocarbon propellants, such as described in U.S. Patent No. 5,376,359 to Johnson.

10 Compositions containing respirable dry particles of micronized active compound of the present invention may be prepared by grinding the dry active compound with, e.g., a mortar and pestle or other appropriate grinding device, and then passing the micronized composition through a 400 mesh screen to break up or separate out large agglomerates.

15 The aerosol, whether formed from solid or liquid particles, may be produced by the aerosol generator at a rate of from about 10 to 150 liters per minute. Aerosols containing greater amounts of medicament may be administered more rapidly. Typically, each aerosol may be delivered to the patient for a period from about 30 seconds to about 20 minutes, with a delivery period of about five to ten minutes being
20 preferred.

The particulate composition comprising the active compound may optionally contain a carrier which serves to facilitate the formation of an aerosol. A suitable carrier is lactose, which may be blended with the active compound in any suitable ratio. Again, other therapeutic compounds such as amiloride, benzamil or phenamil
25 may also be included.

If desired, the active compounds of the present invention may be concurrently administered with uridine 5'-triphosphate (UTP) or an analog thereof (including the pharmaceutically acceptable salts thereof), in an amount effective to stimulate chloride secretion from respiratory epithelial cells (and thereby further hydrate the
30 lung mucus secretions). Formulations containing amiloride, benzamil or phenamil may also contain UTP or an analog thereof in an amount effective to stimulate chloride secretion from respiratory epithelial cells. UTP and analogs thereof that may

be used to carry out this technique are disclosed in U.S. Patent No. 5,292,498 to Boucher.

The present invention is explained in greater detail in the Examples which follow. These examples are intended as illustrative of the invention, and are not to be taken as limiting thereof. In the following Examples, UTP and ATP were obtained from Pharmacia (Uppsala, Sweden), and UDP and hexokinase were from Boehringer Mannheim (Indianapolis, IN). [^3H]myo-inositol (20 Ci/mmol) was from ARC (St. Louis, MO) and [^3H]UTP and [^3H]ATP (97% and 99% pure, respectively) (17-20 Ci/mmol) were from Amersham (Arlington Heights, IL). The miniature perfusion chamber described herein was graciously provided by the staff of E. Larsen's laboratory (Zoophysiological Laboratory A, August Krogh Institute, University of Copenhagen, Denmark). Abbreviations used in the following Examples are as follows: $^{\circ}\text{C}$ means degrees in Centigrade; h means hours; min means minutes; sec means seconds, nm means nanometers; g means grams, ng means nanograms, mg means milligrams, L means liter, mL means milliliter, mmole means millimoles, μmole means micromoles, Ci means curies, μCi means microcuries; and DMEM means Dulbecco's Modified Eagle's Medium.

EXAMPLE 1

Methods: Cell cultures

Human nasal epithelial cells were harvested from turbinates using protease XIV (Sigma, St. Louis, MO) for 24-48 h at 4°C , as previously described. R. Wu, et al., *Am. Rev. Respir. Dis.* **132**, 311-320 (1985). Cytosolic Ca^{2+} and inositol phosphate measurements were made on nasal cell monolayers plated on porous Transwell Col filters (pore diameter $0.45\ \mu\text{m}$; Costar, Cambridge, MA) and maintained in Ham's F12 media supplemented with 10 ng/mL epidermal growth factor, 3.75 ng/mL endothelial cell growth factor, 500 ng/mL hydrocortisone, 5 ng/mL insulin, and 1 mM CaCl_2 (Ham's F12 + 4X medium). Assays were carried out 7 to 10 days after seeding, a time coincident with the development of the maximal transepithelial potential difference according to N. Nakahata & T. K. Harden, *Biochem. J.* **241**, 337-344 (1987).

EXAMPLE 2**Methods: Measurement of inositol phosphates**

5 Confluent human nasal epithelial cells as described in Example 1 were labeled for 18 h in inositol-free DMEM containing 4.5g/L glucose and 5 $\mu\text{Ci/mL}$ of [^3H]myo-inositol. These cells were pre-incubated with 10 mM LiCl and challenged with agonists for an additional 10 min. No changes of medium were made subsequent to addition of [^3H]myo-inositol to avoid release of endogenous
10 ATP from stressed cells. E. R. Lazarowski et al., *Br. J. Pharmacol.* 116, 1619-1627 (1995). Incubations were terminated by addition of 5% ice-cold trichloroacetic acid and the resultant [^3H]inositol phosphates were separated on Dowex AG1-X8 columns as described in A. M. Paradiso et al., *Nature* 377, 643-646 (1995).

15

EXAMPLE 3**Methods: Combined bioelectric and cytosolic Ca^{2+} measurements**

Nasal cells grown on Transwell Col filters affixed to O-rings were
20 maintained in F12 + 4X medium and studied 7-10 days after cell plating as described above. For Ca^{2+}_i measurements, cells were loaded with Fura-2 and mounted in a miniature Ussing chamber over an objective of a microscope (Zeiss) coupled to a microfluorimeter, and cytosolic Ca^{2+}_i levels were quantitated as previously described in Paradiso et al., *supra*. The fluorescence intensity ratio
25 (excitation 340/380; emission λ 450 nm) was collected from a field of 30-40 nasal cells on monolayers and converted to Ca^{2+} as previously described in E. H. Larsen et al., *J. Physiol.* 424, 109-131 (1990). For simultaneous measurements of Cl^- secretion, the transepithelial potential difference (TEP) was measured by a Voltage-Clamp/Pulse Generator (Model VCC600, Physiologic Instruments, San
30 Diego, CA) and recorded on a two-channel recorder (Linsesis Model L200S). To calculate changes in Cl^- secretory current (ΔI_{Cl^-}), a defined 1 sec current pulse was delivered across the monolayer every 10 sec. The nasal tissue was converted

from its native Na⁺ absorptive state to a Cl⁻ secretory state by exposing monolayers to a luminal medium containing 0 Na⁺/low Cl⁻ and to a basolateral medium of Krebs bicarbonate Ringer solution.

5

EXAMPLE 4**Methods: Hydrolysis of [³H]UTP and [³H]UDP**

Nasal monolayers were grown to confluence on 6.5 mm Transwells (Costar) as indicated above. The cells were washed twice and preincubated with 300 µl HEPES-buffered (pH, 7.4) DMEM medium for at least one hour prior to the addition of the [³H]-nucleotides to ensure that any endogeneously released ATP was degraded. Incubations were initiated by addition of 2 µM [³H]UTP or [³H]UDP (0.5 µCi each) to either cell surface. Incubations were terminated at the indicated times by transferring the medium to a tube containing 30 µl 50 mM EDTA and boiled for 2 min.

15

EXAMPLE 5**Methods: Nucleoside diphosphokinase (NDPK) assay**

To assay for the presence of ecto-NDPK activity, cells were incubated with [³H]UDP as detailed above and ATP (100 µM) was included in the incubation medium. The ATP-dependent conversion of [³H]UDP to [³H]UTP was quantitated as described previously in E. R. Lazarowski et al., *Br. J. Pharmacol.* 117, 203-209 (1995).

25

EXAMPLE 6**Methods: HPLC analysis**

Nucleotides were separated by HPLC (Shimadzu Scientific Instruments, Inc., Columbia, MD) via a strong anion exchange column (Rainin Instrument Co., Emeryville, CA) using a two solvent system consisting of buffer A (45 mM ammonium formate, pH 4.6) and buffer B (250 mM sodium phosphate, pH 2.7). A linear gradient was developed from 100% buffer A to 100% B during the first 25 min. The column then was eluted with 100% buffer B for the following 15 min

30

and with 100% buffer A for an additional 15 min from 45 to 60 min. Absorbance at 260 nm was monitored with a LSA UV detector (Shimadzu), and radioactivity was monitored on-line with a Flo-One detector (Packard Instrument Co., Meriden, CT). [^3H]Nucleotides and [^3H]nucleosides were quantitated as previously described. *Id.*

EXAMPLE 7

Methods: Enzymatic conversion of UTP to UDP

Accurate delineation of the pharmacological selectivities of nucleotide receptors is hampered by impurities frequently present in nucleotide solutions and by potential ecto-enzyme-catalyzed interconversion between triphospho- and diphosphonucleotides. Commercial preparations of UDP contain up to 2% UTP. Experiments were performed to evaluate the efficacy of hexokinase in converting UTP to UDP. It is known that in the presence of glucose, hexokinase transfers the γ -phosphate of ATP to glucose, producing ADP and glucose 6-phosphate as products. E. A. Barnard, *Meth. Enzymol.* 42, 6-20 (1975). Although ATP is the preferential substrate for hexokinase, other nucleoside triphosphates also serve as γ -phosphate donors although at slower rates of reaction. *Id.*

It was determined that incubation of 1 mM UTP with 10 U/mL hexokinase results in quantitative conversion of UTP to UDP. Incubations were carried out at 37°C in 1 mL HEPES buffered DMEM medium (pH 7.4) containing 25 mM glucose and 1 mM (0.5 μCi) of either [^3H]ATP or [^3H]UTP. At the times indicated, 5 mM EDTA was added to the samples, followed by boiling. The results are expressed as the percent conversion of [^3H]nucleotide triphosphate to [^3H]nucleotide diphosphate. The data indicate the mean value of one experiment representative of two independent experiments performed with duplicate samples which differed by less than 20%.

In view of these results, and to ensure that no UTP contaminated UDP, stock solutions of UDP (1 mM) were routinely pre-incubated with 10 U/mL hexokinase and 25 mM glucose for 1 h prior to assays.

EXAMPLE 8
Methods: Synthesis of [³H]UDP

[³H]UDP was prepared by incubating [³H]UTP with hexokinase and
5 glucose followed by boiling for 2 min as indicated above.

EXAMPLE 9
Metabolism of [³H]UDP and [³H]UTP by
human airway epithelial cell surfaces

10 Confluent polarized cells were incubated for 20 min at 37°C in the presence
of 1 μM (0.2 μCi) of [³H]UTP, [³H]UDP, or [³H]UDP combined with 100 μM
ATP. All additions were to the mucosal bath (300 μl final volume). Incubations
were terminated by transferring the mucosal medium to an Eppendorf tube
15 containing 30 μl 50 mM EDTA, followed by boiling. [³H]Species were separated
by HPLC as indicated in Example 6.

The results of this experiment are illustrated in the HPLC traces of **FIG.**
2A (incubation with [³H]UTP), **FIG. 2B** (incubation with [³H]UDP), and **FIG.**
2C (incubation with [³H]UDP combined with 100 μM ATP). Each trace is
20 representative of at least three independent experiments performed with duplicates.
Incubation of 1 μM [³H]UTP for 20 min on the mucosal surface resulted in 73 ±
16% hydrolysis, indicating the presence of ecto-nucleotidases and/or phosphatases.
[³H]UDP accumulated as the major breakdown product of [³H]UTP, as shown in
FIG. 2A. Mucosal [³H]UDP (1 μM) was also hydrolyzed, although to a lesser
25 extent than [³H]UTP, as shown in **FIG. 2B**. Time course experiments indicated
that the half life (t_{1/2}) values for mucosal [³H]UTP and [³]UDP were 14 ± 2 min and
27 ± 3 min, respectively (not shown).

Hydrolysis of both [³H]UTP and [³H]UDP on the surface of human nasal
epithelial cells was determined on both cell surfaces by the following procedures:
30 primary cultures of HNE cells were grown to confluence on 6.5 mm Transwells as
a polarized epithelium. The cells were washed twice with pre-warmed DMEM-
HEPES medium (pH 7.4), and were pre-incubated for 1 hour at 37°C with 0.5 mL
of medium added on each side. Incubations were initiated by the addition 50 μl of
10 μM (0.5 μCi) [³H]UTP or [³H]UDP on the indicated cell surface. Medium

samples were collected during a 20 min incubation period and analyzed by HPLC as described in Example 6. The results from these experiments are summarized in **Table 1** below. The values represent the mean (\pm range from the mean) from two experiments performed with duplicate samples.

	Nucleotide Hydrolysis	
	<u>MUCOSAL</u>	<u>SEROSAL</u>
[³ H]UTP	17 \pm 22	98 \pm 11
[³ H]UDP	87 \pm 6	70 \pm 9

5 Nucleoside diphosphokinase catalyzes the transfer of the γ -phosphate from nucleoside triphosphates to nucleoside diphosphates. It has been observed previously that the presence of a nucleoside diphosphokinase activity on the surface of 1321N1 human astrocytoma cells converts UDP (and ADP) to UTP (or ATP), and confounds analyses of the pharmacological effects of

10 diphosphonucleotides. To examine the potential presence of ecto-nucleoside diphosphokinase activity in airway epithelial cells, [³H]UDP was added to the mucosal surface in combination with ATP. These results are shown in **FIG. 2C**. [³H]UTP was rapidly formed under these conditions, and similar results were

15 obtained with either 0.1 μ M or 100 μ M [³H]UDP. Comparable results also were obtained in experiments examining nucleoside diphosphokinase activity on the serosal side. Since potential ATP release from epithelial cells during tissue manipulations and consequential phosphorylation of UDP to UTP by nucleoside diphosphokinase may complicate study of the actions of UDP, hexokinase (2 U/mL) and glucose were included in all subsequent assays examining the effects of

20 UDP, as explained above. No accumulation of [³H]UTP occurred under these incubation conditions.

EXAMPLE 10

Activity of UDP at Human Nasal Epithelial Cell Surface

25 It has been established that primary cultures of polarized human airway epithelial cells express P2Y₂ receptors on both cell surfaces. S. J. Mason et al., *Br. J. Pharmacol.* 103, 1649-1656 (1991). It has also been reported that functional

expression of the P2Y₂ receptor in polarized human nasal epithelial cells is asymmetric and that the receptor apparently couples more efficaciously to its effector phospholipase C on the serosal (basolateral) surface than on the mucosal (apical) surface. Paradiso et al., *supra*.

5 Consistent with this previous study, UTP (100 μ M) promoted larger [³H]inositol phosphate and calcium responses when applied to the basolateral bath of polarized primary human nasal epithelial cells, than when applied to the mucosal bath, as shown in **FIGS. 3A and 3B**. The results of **FIGS. 3A and 3B** were obtained after confluent cells were loaded with [³H]*myo*-inositol and
10 preincubated with LiCl (**FIG. 3A**, or with Fura-2 (**FIG. 3B**), as described above in Examples 2 and 3. The cells were challenged with 100 μ M of the indicated nucleotide, added to either the serosal or the mucosal medium. The Cl⁻ secretory responses (ΔI_{Cl^-}) following addition of 100 μ M UTP to either the mucosal or the serosal bath were $74 \pm 11 \mu A/cm^2$ and $34 \pm 3 \mu A/cm^2$, respectively. Application
15 of UDP to the serosal bath had negligible effect on inositol phosphate accumulation.

In contrast, mucosal UDP (100 μ M) promoted marked accumulation of [³H]inositol phosphates and calcium mobilization, and the maximal effects of UDP were approximately one half the magnitude of the responses observed with
20 mucosal UTP, as shown in **FIGS. 3 and 4**. Similarly, mucosal but not serosal UDP stimulated Cl⁻ secretion ($\Delta I_{Cl^-} = 16 \pm 3 \mu A \cdot cm^2$ for 3 μ M mucosal UDP; $\Delta I_{Cl^-} = 0 \pm 0 \mu A \cdot cm^2$ for 3 μ M serosal UDP) and the magnitude of this response was approximately one half of the mucosal UTP response (shown in **FIG. 4**). The effects of mucosal application of UDP cannot be explained by activation of P2Y₂
25 receptors since UTP-free UDP is essentially inactive at the P2Y₂ receptor (Nicholas et al., *supra*), and since UDP caused little or no effect when applied to the P2Y₂ receptor-expressing serosal side of the cell monolayers (**FIGS. 3 and 4**). The results in **FIG. 4** were obtained after primary cultures of human nasal epithelial cells were mounted on modified Ussing chambers as detailed above in
30 Example 3. Changes in Ca²⁺ concentration (top tracings) or in Cl⁻ secretion

(bottom tracings) were simultaneously recorded after the addition to the basolateral medium of 100 μ M UDP followed by 100 μ M UTP (serosal), or after two consecutive additions of 3 μ M UDP and then 10 μ M UDP to the apical medium followed by the ipsilateral addition of 10 μ M UTP (mucosal). The larger mucosal versus serosal effect of UDP contrasts with the predominantly basolateral effects of UTP (FIG. 3) and ATP. *See, e.g.,* Paradiso et al., *supra.* Thus, the action of UDP in polarized airway epithelial cells does not coincide with that observed with P2Y₂ receptor agonists.

Although mucosal UDP was less efficacious than mucosal UTP in stimulating inositol phosphate formation in human nasal epithelium, UDP (EC₅₀ = 190 \pm 27 nM) and UTP (EC₅₀ = 280 \pm 35 nM) exhibited similar potencies, as shown in FIG. 5. The results in FIG. 5 were obtained after confluent cells were labeled with [³H]myo-inositol, preincubated with LiCl as described above, and subsequently challenged for 10 min with the indicated concentration of UDP or UTP added to the mucosal medium. Cross-desensitization experiments were carried out to further address the hypothesis that UDP promotes signaling responses in airway cells through a receptor distinct from the P2Y₂ receptor. Addition of 3 μ M UDP to the mucosal bath promoted mobilization of intracellular calcium with a Δ Ca²⁺ of 66 \pm 12 nM, and changes in Cl⁻ secretory responses (Δ I_{Cl⁻) of -16.3 \pm 3 μ A/cm² (n = 8). Subsequent addition of 3 μ M UDP followed by 10 μ M UDP did not result in elevation of intracellular Ca²⁺ (Δ Ca²⁺ = 0) or in Cl⁻ secretion (Δ I_{Cl⁻ = 0), which suggests the occurrence of UDP-induced desensitization (see FIG. 4). In contrast, responses to UTP (control cells, Δ Ca²⁺ = 374 \pm 24 nM [n = 19]; Δ I_{Cl⁻ = -74 \pm 11 μ A/cm² [n = 12]; UDP-treated cells, Δ Ca²⁺ = 310 \pm 35 nM [n = 8]; Δ I_{Cl⁻ = -61 \pm 10 nM [n = 8]) were retained. Responses to ATP also were retained following multiple additions of UDP.}}}}

EXAMPLE 11
Analysis of UDP Assay Results

The results presented above illustrate that although UDP is not a P2Y₂ receptor agonist, it nonetheless is a potent agonist at the human airway epithelial cell surface. The effects of UDP could not be attributed to contamination with UTP. Solutions of UDP were pre-incubated with hexokinase and glucose to eliminate any traces of UTP, and hexokinase and glucose were included in studies of epithelial cells to prevent metabolic conversion of UDP to UTP by cell surface nucleoside diphosphokinases and endogenously released ATP. HPLC analysis of UDP solutions before and after incubation with cells indicated the absence of UTP. The almost equal potency of UDP and UTP on the airway cells notably contrasts with the low apparent activity of UDP at P2Y₂ receptors. The asymmetry of the effect of UDP on polarized epithelium, with a preferred mucosal versus serosal activity, also contrasts with the predominant serosal effect of UTP (and ATP) in airway cells. Since UDP does not activate the epithelial cell P2Y₂ receptor, the explanation for the stimulatory effect of this diphosphate is the existence of a novel airway receptor that selectively recognizes UDP.

The potential implications of these results are two-fold. First, they demonstrate that in the presence of the apparently more abundant P2Y₂ receptor, assay conditions can be established that allow resolution of a receptor that is selectively activated by UDP. Specifically, hexokinase and glucose can be utilized to study effects of UDP in tissues where UDP-selective effects are otherwise masked by a prominent P2Y₂ receptor. Second, the asymmetry of the effects of UDP and UTP in polarized epithelial cells indicates different regulatory functions for the receptors recognizing these two nucleotides. Activators of receptors on the serosal surface may be released at distant sites and access receptors via the blood stream, whereas UDP, which acts exclusively on the mucosal side of airway epithelial cells, may be generated locally. However, mucosal UTP is hydrolyzed at a 2-3 fold faster rate than mucosal UDP, and consequently, UDP likely accumulates on the apical surface after the breakdown of UTP. Thus, released

UTP could be an important source of extracellular UDP, which in turn potentially serves as a physiologically important signaling molecule in human airway tissue.

EXAMPLE 12

Effect of UDP on mucociliary clearance in sheep

5 Healthy adult ewes were given ^{99m}Tc -labeled serum albumin (^{99m}Tc -HCA) via a nebulized aerosol. The (^{99m}Tc -HCA)(20 mCi) was administered over 5 minutes through a nasotracheal tube introduced under local anesthesia with 2%
10 lidocaine. After administration of the ^{99m}Tc -HCA, the animals were given the test compound UDP. The test compound was administered by nebulization in a volume of 4 mL over a period of 10-12 min. The test compound was given at a dose of 400 μmole . After the administration of the test compound, the animals were extubated. Clearance of the radiolabeled particles was monitored with a
15 gamma camera. Measurements were made at 0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 75, 90, 105 and 120 minutes. Results of this study ($n=7$) have shown that the test compound promotes clearance of radiolabeled particles when compared to the saline control. The maximal effect for 400 μmole UDP was $33.6 \pm 1.6\%$ clearance. The response to UDP was maximal at 60 minutes and was
20 maintained for the second hour of the study. The results of this experiment are illustrated graphically in FIG. 6.

EXAMPLE 13

Effect of UDP on tracheal mucus velocity in sheep

25 To measure the effect of UDP on tracheal mucus velocity (TMV), the nasal passages of conscious adult ewes were anesthetized with a 2% lidocaine solution. After local anesthesia was produced, a modified endotracheal tube (7.5mm) was placed such that the cuff was just below the vocal cords (as verified by
30 fluoroscopy). Inspired air was warmed and humidified. The cuff on the endotracheal tube was inflated only during administration of the test compound to minimize possible impairment of TMV by the cuff. Test compounds were administered by nebulization in a volume of 4 mL over a period of 10-12 min.

TMV was measured by fluoroscopy. Ten to twenty radiopaque disks (Teflon/bismuth trioxide; 1mm diameter; 0.8mm thick, weighing 1.8 mg) were introduced into the trachea through a modified suction catheter with a puff of compressed air (3-4 L/min). Velocities of the individual disks were recorded on videotape from a portable image intensifier unit. Individual disk velocities were calculated by measuring the distance traveled by each disk during a 1 min observation period. Values reported are the means of the individual disk velocities. A collar was worn by the sheep, which was used as a standard to correct for magnification errors inherent in the fluoroscope.

UDP (400 μ mole; 4 mL of a 10^{-1} M) produced significant effects on tracheal mucus velocity. UDP produced a maximal effect of $121 \pm 8\%$ of baseline (mean standard error, n=6). UDP produced its maximal effects 15 min after administration. However, when pairwise comparisons were made, only the time point immediately after dosing (t=0) was significantly different from the saline treatment group. The results of this experiment are illustrated graphically in FIG. 7.

EXAMPLE 14

Synthesis of UDP Analogs and P2Y₆ Activity of Analogs

The following method was used for measuring inositol phosphate accumulation of the UDP analogs described within this example as drugs. 1321N1 astrocytoma cells, stably overexpressing the human P2Y₆ receptor (Nicholas, et al., *Mol. Pharmacol.* 50, 224-229 (1996)), were seeded into 96 well plates (7500 cells/well) in DMEM-H supplemented with 10% FCS. After 48 hours, the growth media was replaced with inositol-free DMEM-H supplemented with 2.5% dialyzed FCS and [³H]-inositol (0.2 μ Ci/well) to radiolabel hormone-responsive inositol phospholipid pools. Cells were stimulated 48 hours later. At the time of assay, the culture media was supplemented with 25 mM HEPES (pH 7.4) and 10 mM LiCl₂ to prevent hydrolysis of [³H]-inositol phosphates. In addition, hexokinase (0.03 U/well) and glucose (50 mM final concentration) was added to the cultures to provide an enzymatic mechanism for the depletion of nucleoside triphosphates contaminating drug solutions or released from cells prior to/or during cell

stimulation reactions. Drugs were diluted in phosphate buffered saline containing hexokinase/glucose and incubated at 37° for 30 minutes prior to addition to cells. Reactions were terminated after 60 minutes by rapid aspiration of media followed by the addition of ice cold EDTA (1.0 mM) and incubation on ice for at least 15
5 minutes prior to resolution of [³H]-inositol phosphates by ion exchange chromatography as described (Brown, et. al., *supra*).

The drugs used in the assay described above were synthesized as follows:

(Compound A) 5-Methyluridine 5'-diphosphate

5-Methyluridine 5'-triphosphate (0.0047g, 0.009mMol) was dissolved in
10 0.7 mL 1M Tris buffer (pH 8.5 containing 1M MgCl₂) then 0.3 mL of a 0.25M glucose solution in 1M Tris buffer (pH 8.5 containing 1M MgCl₂) added. A T₀ timepoint was taken by reverse phase anion exchange HPLC. Hexokinase (EC 2.7.1.1, Boehringer Mannheim, from yeast overproducer), 75 units in 3.2M ammonium sulfate suspension, was added to the reaction which was incubated at
15 25°C. An HPLC was taken at 66 hours indicating complete conversion to diphosphate with no triphosphate remaining. Triphosphate retention: 17.5 minutes; diphosphate retention: 10.3 minutes.

The following nucleoside 5'-diphosphates were prepared by the same procedure as above, with modifications noted.

20

(Compound B) 4-Hexyloxyuridine 5'-diphosphate

Incubation for 21 days, diphosphate retention: 13.5 minutes.

(Compound C) N,N-dimethylcytidine 5'-diphosphate

25 Incubation for 30 days, diphosphate retention: 11.62 minutes.

(Compound D) 4-Methoxyuridine 5'-diphosphate

Incubation for 20 days, diphosphate retention: 10.76 minutes.

30 **(Compound E) N-hexylcytidine 5'-diphosphate**

Incubation for 12 days, diphosphate retention: 11.56 minutes.

(Compound F) 4-(N-morpholino)uridine 5'-diphosphate

Incubation for 21 days, diphosphate retention: 10.78 minutes.

(Compound G) 5-(Phenylethynyl)uridine 5'-diphosphate

5 Incubation for 48 hours, diphosphate retention: 13.91 minutes.

(Compound H) 3'-Deoxyuridine 5'-diphosphate

Incubation for 35 days, diphosphate retention: 8.0 minutes.

10 (Compound I) N-Cyclopentylcytidine 5'-diphosphate

Incubation for 6 days, diphosphate retention: 12.24 minutes.

(Compound J) 4-Hexylthiouridine 5'-diphosphate

Incubation for 4 days, diphosphate retention: 7.51 minutes.

15

Activity on P2Y₆ was measured as described above, with the following

results:

Activity on P2Y₆ (EC₅₀ in μ Mol)

	Compound A:	0.5
20	Compound B:	3.0
	Compound C:	41.7
	Compound D:	4.7
	Compound E:	25.5
	Compound F:	55.1
25	Compound G:	0.02
	Compound H:	2.1
	Compound I:	8.6
	Compound J:	1.8


30

The foregoing Examples are illustrative of the present invention and are not to be construed as limiting thereof. Accordingly, the invention is defined by the following claims, with equivalents of the claims included therein.

THAT WHICH IS CLAIMED:

1. A method of hydrating lung mucus secretions in the lungs of a subject in need of such treatment, comprising administering to the lungs of the subject a compound of Formula I below, or a pharmaceutically acceptable salt thereof, in an amount sufficient to hydrate lung mucus secretions:

5



15 wherein:

X_1 , and X_2 are each independently either O^- or S^- ;

X_3 and X_4 are each independently either $-H$ or $-OH$, with the proviso that X_3 and X_4 are not simultaneously $-H$;

R_1 is selected from the group consisting of O , imido, methylene and dihalomethylene;

R_2 is selected from the group consisting of H , halo, alkyl, substituted alkyl, alkoxyl, nitro and azido;

R_3 is selected from the group consisting of H , alkyl, acyl, aryl, and arylalkyl; and

25 R_4 is selected from the group consisting of $-OR'$, $-SR'$, $-NR'$, and $-NR'R''$, wherein R' and R'' are independently selected from the group consisting of H , alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxyl, and aryloxyl, and with the proviso that R' is absent when R_4 is double bonded from an oxygen or sulfur atom to the carbon at the 4-position of the pyrimidine ring.


2. A method according to claim 1, wherein said compound of Formula I is administered by delivering an aerosol suspension of respirable particles comprised of a compound of Formula I to the lungs of said subject.

3. A method according to claim 2, wherein said particles are selected from the group consisting of solid particles and liquid particles.

4. A method according to claim 2, wherein said aerosol is comprised of particles in the respirable range.

5. A method of treating cystic fibrosis in a human subject in need of such treatment, comprising administering by inhalation an aerosol suspension of respirable particles to the lungs of said subject, said particles comprised of Formula I, or a pharmaceutically acceptable salt thereof :

5



10

wherein:

15

X_1 , and X_2 are each independently either O^- or S^- ;

X_3 and X_4 are each independently either $-H$ or $-OH$, with the proviso that X_3 and X_4 are not simultaneously $-H$;

R_1 is selected from the group consisting of O , imido, methylene and dihalomethylene;

R₂ is selected from the group consisting of H, halo, alkyl, substituted alkyl, alkoxy, nitro and azido;

R₃ is selected from the group consisting of H, alkyl, acyl, aryl, and arylalkyl; and

5 R₄ is selected from the group consisting of —OR', —SR', —NR', and —NR'R", wherein R' and R" are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxy, and aryloxy, and with the proviso that R' is absent when R₄ is double bonded from an oxygen or sulfur atom to the carbon at the 4-position of the pyrimidine ring;

10 and wherein said particles are administered in an amount effective to treat said cystic fibrosis in said subject.


6. A method according to claim 5, wherein said particles are selected from the group consisting of solid particles and liquid particles.

7. A method according to claim 6, wherein said aerosol is comprised of particles having a particle size within the respirable range.

8. A method according to claim 5, further comprising concurrently administering to said subject a compound selected from the group consisting of amiloride, benzamil, and phenamil.

9. A method of treating primary ciliary dyskinesia in a human subject in need of such treatment, comprising administering by inhalation an aerosol suspension of respirable particles to the lungs of said subject, said particles comprised of Formula

I, or a pharmaceutically acceptable salt thereof :



wherein:

X_1 , and X_2 are each independently either O^- or S^- ;

15 X_3 and X_4 are each independently either $-H$ or $-OH$, with the proviso that X_3 and X_4 are not simultaneously $-H$;

R_1 is selected from the group consisting of O , imido, methylene and dihalomethylene;

R_2 is selected from the group consisting of H , halo, alkyl, substituted alkyl, alkoxy, nitro and azido;

20 R_3 is selected from the group consisting of H , alkyl, acyl, aryl, and arylalkyl; and

R_4 is selected from the group consisting of $-OR'$, $-SR'$, $-NR'$, and $-NR'R''$, wherein R' and R'' are independently selected from the group consisting of H , alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxy, and aryloxy, and with the proviso that R' is absent when R_4 is double bonded from an oxygen or sulfur atom to the carbon at the 4-position of the pyrimidine ring;

25 and wherein said particles are administered in an amount effective to treat said primary ciliary dyskinesia in said subject.

10. A method according to claim 9, wherein said particles are selected from the group consisting of solid particles and liquid particles.


11. A method according to claim 10, wherein said aerosol is comprised of particles having a particle size within the respirable range.

12. A method according to claim 9, further comprising concurrently administering to said subject a compound selected from the group consisting of amiloride, benzamil, and phenamil.

13. A method of treating chronic obstructive pulmonary disorder in a human subject in need of such treatment, comprising administering by inhalation an aerosol suspension of respirable particles to the lungs of said subject, said particles comprised of Formula I, or a pharmaceutically acceptable salt thereof :

5

10



15 wherein:

X_1 , and X_2 are each independently either O^- or S^- ;

X_3 and X_4 are each independently either $-H$ or $-OH$, with the proviso that X_3 and X_4 are not simultaneously $-H$;

R_1 is selected from the group consisting of O , imido, methylene and dihalomethylene;

R_2 is selected from the group consisting of H , halo, alkyl, substituted alkyl, alkoxy, nitro and azido;

R₃ is selected from the group consisting of H, alkyl, acyl, aryl, and arylalkyl;
and

R₄ is selected from the group consisting of —OR', —SR', —NR', and
—NR'R", wherein R' and R" are independently selected from the group consisting of
5 H, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxyl, and aryloxy, and
with the proviso that R' is absent when R₄ is double bonded from an oxygen or sulfur
atom to the carbon at the 4-position of the pyrimidine ring;

and wherein said particles are administered in an amount effective to treat
said chronic obstructive pulmonary disorder in said subject.


14. A method according to claim 13, wherein said particles are selected from
the group consisting of solid particles and liquid particles.

15. A method according to claim 14, wherein said aerosol is comprised of
particles having a particle size within the respirable range.

16. A method according to claim 13, further comprising concurrently
administering to said subject a compound selected from the group consisting of
amiloride, benzamil, and phenamil.

17. A method of treating ventilator-associated pneumonia in a human subject
in need of such treatment, comprising administering by inhalation an aerosol
suspension of respirable particles to the lungs of said subject, said particles comprised

of Formula I, or a pharmaceutically acceptable salt thereof :



wherein:

X_1 , and X_2 are each independently either O^- or S^- ;

X_3 and X_4 are each independently either $-H$ or $-OH$, with the proviso that X_3 and X_4 are not simultaneously $-H$;

15 R_1 is selected from the group consisting of O , imido, methylene and dihalomethylene;

R_2 is selected from the group consisting of H , halo, alkyl, substituted alkyl, alkoxyl, nitro and azido;

R_3 is selected from the group consisting of H , alkyl, acyl, aryl, and arylalkyl;

20 and

R_4 is selected from the group consisting of $-OR'$, $-SR'$, $-NR'$, and $-NR'R''$, wherein R' and R'' are independently selected from the group consisting of H , alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxyl, and aryloxyl, and with the proviso that R' is absent when R_4 is double bonded from an oxygen or sulfur atom to the carbon at the 4-position of the pyrimidine ring;


25 and wherein said particles are administered in an amount effective to treat said ventilator-associated pneumonia in said subject.

18. A method according to claim 17, wherein said particles are selected from the group consisting of solid particles and liquid particles.

19. A method according to claim 18, wherein said aerosol is comprised of particles having a particle size within the respirable range.

20. A method according to claim 17, further comprising concurrently administering to said subject a compound selected from the group consisting of amiloride, benzamil, and phenamil.

21. A method of treating chronic bronchitis in a human subject in need of such treatment, comprising administering by inhalation an aerosol suspension of respirable particles to the lungs of said subject, said particles comprised of Formula I, or a pharmaceutically acceptable salt thereof:



wherein:

X_1 , and X_2 are each independently either O^- or S^- ;

15 X_3 and X_4 are each independently either $-H$ or $-OH$, with the proviso that X_3 and X_4 are not simultaneously $-H$;

R_1 is selected from the group consisting of O , imido, methylene and dihalomethylene;

20 R_2 is selected from the group consisting of H , halo, alkyl, substituted alkyl, alkoxy, nitro and azido;

R_3 is selected from the group consisting of H , alkyl, acyl, aryl, and arylalkyl;

and

R_4 is selected from the group consisting of $-OR'$, $-SR'$, $-NR'$, and $-NR'R''$, wherein R' and R'' are independently selected from the group consisting of

25 H , alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxy, and aryloxy, and

with the proviso that R' is absent when R₄ is double bonded from an oxygen or sulfur atom to the carbon at the 4-position of the pyrimidine ring;


and wherein said particles are administered in an amount effective to treat said chronic bronchitis in said subject.

22. A method according to claim 21, wherein said particles are selected from the group consisting of solid particles and liquid particles.

23. A method according to claim 22, wherein said aerosol is comprised of particles having a particle size within the respirable range.

24. A method according to claim 21, further comprising concurrently administering to said subject a compound selected from the group consisting of amiloride, benzamil, and phenamil.

25. A pharmaceutical composition, comprising, together in a pharmaceutically acceptable carrier, a compound of Formula I:



wherein:

15 X_1 , and X_2 are each independently either O^- or S^- ;

X_3 and X_4 are each independently either $-H$ or $-OH$, with the proviso that X_3 and X_4 are not simultaneously $-H$;

R₁ is selected from the group consisting of O, imido, methylene and dihalomethylene;

R₂ is selected from the group consisting of H, halo, alkyl, substituted alkyl, nitro and azido;

5 R₃ is selected from the group consisting of H, alkyl, acyl, aryl, and arylalkyl; and

R₄ is selected from the group consisting of —OR', —SR', —NR', and —NR'R'', wherein R' and R'' are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxy, and aryloxy, and
10 with the proviso that R' is absent when R₄ is double bonded from an oxygen or sulfur atom to the carbon at the 4-position of the pyrimidine ring;

or a pharmaceutically acceptable salt thereof, in an amount effective to hydrate lung mucus secretions.


26. A pharmaceutical composition according to claim 25 wherein said carrier is selected from the group consisting of solid carriers and liquid carriers.

27. A pharmaceutical composition according to claim 25 wherein said compound of Formula (I) is selected from the group consisting of uridine 5'-diphosphate; uridine 5'-O-(2-thiodiphosphate); 2-deoxyuridine 5'-diphosphate; and 4-mercaptouridine 5'-diphosphate, and the pharmaceutically acceptable salts thereof.

28. A pharmaceutical composition according to claim 25 wherein said compound of Formula (I) is selected from the group consisting of 3'-deoxyuridine-5'-diphosphate; 5-(1-phenylethynyl)-uridine 5'-diphosphate; 5-methyluridine 5'-diphosphate; 4-hexylthiouridine 5'-diphosphate; 4-mercaptouridine 5'-diphosphate;
5 4-methoxyuridine 5'-diphosphate; 4-hexyloxyuridine 5'-diphosphate; N,N-dimethylcytidine 5'-diphosphate; N-hexylcytidine 5'-diphosphate; and the pharmaceutically acceptable salts thereof.

29. A pharmaceutical composition according to claim 25, wherein said composition further comprises a propellant.

30. A compound according to Formula I:



wherein:

X_1 , and X_2 are each independently either O^- or S^- ;

X_3 and X_4 are each independently either $-H$ or $-OH$, with the proviso that X_3 and X_4 are not simultaneously $-H$;

15 R_1 is selected from the group consisting of O , imido, methylene and dihalomethylene;

R_2 is selected from the group consisting of H , halo, alkyl, substituted alkyl, alkoxy, nitro and azido;

R_3 is selected from the group consisting of H , alkyl, acyl, aryl, and arylalkyl;

20 and

R_4 is selected from the group consisting of $-OR'$, $-SR'$, $-NR'$, and $-NR'R''$, wherein R' and R'' are independently selected from the group consisting of H , alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, alkoxy, and aryloxy, and with the proviso that R' is absent when R_4 is double bonded from an oxygen or sulfur atom to the carbon at the 4-position of the pyrimidine ring; and;

25 with the proviso that said compound of Formula I is not a compound selected from the group consisting of uridine 5'-diphosphate, 2-deoxyuridine 5'-diphosphate, uridine 5'-O-(2-thiodiphosphate), and 4-mercaptopuridine 5'-diphosphate.

31. A compound according to claim 30, wherein said compound of Formula I is 3'-deoxyuridine-5'-diphosphate.

32. A compound according to claim 30, wherein said compound of Formula I is 5-(1-phenylethynyl)-uridine 5'-diphosphate .

33. A compound according to claim 30, wherein said compound of Formula I is 5-methyluridine 5'-diphosphate.

34. A compound according to claim 30, wherein said compound of Formula I is 4-hexylthiouridine 5'-diphosphate.

35. A compound according to claim 30, wherein said compound of Formula I is 4-mercaptopuridine 5'-diphosphate.

36. A compound according to claim 30, wherein said compound of Formula I is 4-methoxyuridine 5'-diphosphate.

37. A compound according to claim 30, wherein said compound of Formula I is 4-hexyloxyuridine 5'-diphosphate.

38. A compound according to claim 30, wherein said compound of Formula I is N,N-dimethylcytidine 5'-diphosphate.

39. A compound according to claim 30, wherein said compound of Formula I is N-hexylcytidine 5'-diphosphate.

40. A compound according to claim 30, wherein said compound of Formula I is N-cyclopentylcytidine 5'-diphosphate.

1/6

FIG. 1





FIG. 5



2/6

FIG. 2A





FIG. 2B



FIG. 2C



3/6

FIG. 3A




FIG. 3B






FIG. 4




5/6



6/6

FIG. 7



INTERNATIONAL SEARCH REPORT

Internal Application No

PCT/US 98/17894

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/70 C07H19/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 08593 A (UNIVERSITY OF NORTH CAROLINA) 28 April 1994 ---	
A	WO 92 11016 A (FISONS PLC) 9 July 1992 ---	
A	K.N. OLIVER ET AL.: "Acute safety and effects on mucociliary clearance of aerosolized uridine 5'-triphosphate +/- amiloride in normal human adults" AM. J. RESPIR. CRIT. CARE MED., vol. 154, 1996, pages 217-223, XP002056054 ---	
A	DE 25 30 862 A (CHUGAI SEIYAKU KK) 29 January 1976 -----	

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

*** Special categories of cited documents :**

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "S" document member of the same patent family

Date of the actual completion of the international search

30 November 1998

Date of mailing of the international search report

08/12/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Klaver, T

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/ 17894

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 5-24
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 98/17894

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9408593 A	28-04-1994	US 5292498 A	08-03-1994
		AU 688258 B	12-03-1998
		AU 2872592 A	09-05-1994
		EP 0663830 A	26-07-1995
		JP 8502078 T	05-03-1996
		NO 951340 A	02-06-1994
WO 9211016 A	09-07-1992	NONE	
DE 2530862 A	29-01-1976	FR 2277591 A	06-02-1976
		GB 1512043 A	24-05-1978
		US 4058601 A	15-11-1977